



Come experience the Cuemath methodology and ensure your child stays ahead at math this summer.



Adaptive

Platform





Interactive Visual Simulations

For Grades 1 - 10





LIVE online classes by trained and certified experts.

Get the Cuemath advantage

Book a FREE trial class



NCERT Solutions Class 11 Maths Chapter 8 Binomial Theorem

Question 1:

Expand the expression $(1-2x)^5$

Solution:

By using Binomial Theorem, the expression $(1-2x)^5$ can be expanded as $(1-2x)^5 = {}^5C_0(1)^5 - {}^5C_1(1)^4(2x) + {}^5C_2(1)^3(2x)^2 - {}^5C_3(1)^2(2x)^3 + {}^5C_4(1)(2x)^4 - {}^5C_5(2x)^5$ $= 1-5(2x)+10(4x^2)-10(8x^3)+5(16x^4)-(32x^5)$ $= 1-10x+40x^2-80x^3+80x^4-32x^5$

Question 2:

Expand the expression $\left(\frac{2}{x} - \frac{x}{2}\right)^5$

Solution:

By using Binomial Theorem, the expression
$$\left(\frac{2}{x} - \frac{x}{2}\right)$$
 can be expanded as
 $\left(\frac{2}{x} - \frac{x}{2}\right)^5 = {}^5C_0\left(\frac{2}{x}\right)^5 - {}^5C_1\left(\frac{2}{x}\right)^4 \left(\frac{x}{2}\right) + {}^5C_2\left(\frac{2}{x}\right)^3 \left(\frac{x}{2}\right)^2 - {}^5C_3\left(\frac{2}{x}\right)^2 \left(\frac{x}{2}\right)^3 + {}^5C_4\left(\frac{2}{x}\right) \left(\frac{x}{2}\right)^4 - {}^5C_5\left(\frac{x}{2}\right)^5$
 $= \frac{32}{x^5} - 5\left(\frac{16}{x^4}\right) \left(\frac{x}{2}\right) + 10\left(\frac{8}{x^3}\right) \left(\frac{x^2}{4}\right) - 10\left(\frac{4}{x^2}\right) \left(\frac{x^3}{8}\right) + 5\left(\frac{2}{x}\right) \left(\frac{x^4}{16}\right) - \frac{x^5}{32}$
 $= \frac{32}{x^5} - \frac{40}{x^3} + \frac{20}{x} - 5x + \frac{5}{8}x^3 - \frac{x^5}{32}$

Question 3:

Expand the expression $(2x-3)^6$ Solution:

By using Binomial Theorem, the expression $(2x-3)^6$ can be expanded as

$$(2x-3)^{6} = {}^{6}C_{0}(2x)^{6} - {}^{6}C_{1}(2x)^{5}(3) + {}^{6}C_{2}(2x)^{4}(3)^{2} - {}^{6}C_{3}(2x)^{3}(3)^{3} + {}^{6}C_{4}(2x)^{2}(3)^{4} - {}^{6}C_{5}(2x)(3)^{5} + {}^{6}C_{6}(3)^{6}$$

= $64x^{6} - 6(32x^{5})(3) + 15(16x^{4})(9) - 20(8x^{3})(27) + 15(4x^{2})(81) - 6(2x)(243) + 729$
= $64x^{6} - 576x^{5} + 2160x^{4} - 4320x^{3} + 4860x^{2} - 2916x + 729$

Question 4:

Expand the expression $\left(\frac{x}{3} + \frac{1}{x}\right)^5$



By using Binomial Theorem, the expression $\left(\frac{x}{3} + \frac{1}{x}\right)^5$ can be expanded as

$$\begin{split} \left(\frac{x}{3} + \frac{1}{x}\right)^5 &= {}^5C_0 \left(\frac{x}{3}\right)^5 - {}^5C_1 \left(\frac{x}{3}\right)^4 \left(\frac{1}{x}\right) + {}^5C_2 \left(\frac{x}{3}\right)^3 \left(\frac{1}{x}\right)^2 + {}^5C_3 \left(\frac{x}{3}\right)^2 \left(\frac{1}{x}\right)^3 + {}^5C_4 \left(\frac{x}{3}\right) \left(\frac{1}{x}\right)^4 + {}^5C_5 \left(\frac{1}{x}\right)^5 \\ &= \frac{x^5}{243} + 5 \left(\frac{x}{81}\right) \left(\frac{1}{x}\right) + 10 \left(\frac{x^3}{27}\right) \left(\frac{1}{x^2}\right) + 10 \left(\frac{x^2}{9}\right) \left(\frac{1}{x^3}\right) + 5 \left(\frac{x}{3}\right) \left(\frac{1}{x^4}\right) + \frac{1}{x^5} \\ &= \frac{x^5}{243} + \frac{5x^3}{81} + \frac{10x}{27} + \frac{10}{9x} + \frac{5}{3x^3} + \frac{1}{x^5} \end{split}$$

Question 5:

Expand the expression $\left(x+\frac{1}{x}\right)^6$

Solution:

By using Binomial Theorem, the expression
$$\binom{x+\frac{1}{x}}{x}$$
 can be expanded as
 $\left(x+\frac{1}{x}\right)^6 = {}^6C_0\left(x\right)^6 + {}^6C_1\left(x\right)^5\left(\frac{1}{x}\right) + {}^6C_2\left(x\right)^4\left(\frac{1}{x}\right)^2 + {}^6C_3\left(x\right)^3\left(\frac{1}{x}\right)^3 + {}^6C_4\left(x\right)^2\left(\frac{1}{x}\right)^4 + {}^6C_5\left(x\right)\left(\frac{1}{x}\right)^5 + {}^6C_6\left(\frac{1}{x}\right)^6$
 $= x^6 + 6\left(x\right)^5\left(\frac{1}{x}\right) + 15\left(x\right)^4\left(\frac{1}{x^2}\right) + 20\left(x\right)^3\left(\frac{1}{x^3}\right) + 15\left(x\right)^2\left(\frac{1}{x^4}\right) + 6\left(x\right)\left(\frac{1}{x^5}\right) + \left(\frac{1}{x^6}\right)$
 $= x^6 + 6x^4 + 15x^2 + 20 + \frac{15}{x^2} + \frac{6}{x^4} + \frac{1}{x^6}$

Question 6:

Using Binomial Theorem, evaluate $(96)^3$

Solution:

96 can be expressed as the difference of two numbers whose powers are easier to calculate. Hence, 96 = 100 - 4

Therefore,

$$(96)^{3} = (100 - 4)^{3}$$

$$= {}^{3}C_{0} (100)^{3} - {}^{3}C_{1} (100)^{2} (4) + {}^{3}C_{2} (100) (4)^{2} - {}^{3}C_{3} (4)^{3}$$

$$= (100)^{3} - 3 (10000) (4) + 3 (100) (16) - (64)$$

$$= 1000000 - 120000 + 4800 - 64$$

$$= 1004800 - 120064$$

$$= 884736$$



Question 7:

Using Binomial Theorem, evaluate $(102)^5$

Solution:

102 can be expressed as the sum of two numbers whose powers are easier to calculate. Hence, 102 = 100 + 2Therefore; $(102)^5 = (100 + 2)^5$ $= {}^5C_0 (100)^5 + {}^5C_1 (100)^4 (2) + {}^5C_2 (100)^3 (2)^2 + {}^5C_3 (100)^2 (2)^3 + {}^5C_4 (100) (2)^4 + {}^5C_5 (2)^5$ = 1000000000 + 100000000 + 4000000 + 800000 + 8000 + 32= 11040808032

Question 8:

Using Binomial Theorem, evaluate $(101)^4$

Solution:

101 can be expressed as the sum of two numbers whose powers are easier to calculate. Hence 101 = 100 + 1Therefore,

$$(101)^{4} = (100+1)^{4}$$

= ${}^{4}C_{0} (100)^{4} + {}^{4}C_{1} (100)^{3} (1) + {}^{4}C_{2} (100)^{2} (1)^{2} + {}^{4}C_{3} (100) (1)^{3} + {}^{4}C_{4} (1)^{4}$
= $(100)^{4} + 4 (100)^{3} + 6 (100)^{2} + 4 (100) + (1)^{4}$
= $100000000 + 4000000 + 60000 + 400 + 1$
= 104060401

Question 9:

Using Binomial Theorem, evaluate $(99)^5$

Solution:

99 can be expressed as the difference of two numbers whose powers are easier to calculate.

Hence, 99 = 100 - 1



Therefore,

$$(99)^{5} = (100-1)^{5}$$

= ${}^{5}C_{0}(100)^{5} - {}^{5}C_{1}(100)^{4}(1) + {}^{5}C_{2}(100)^{3}(1)^{2} - {}^{5}C_{3}(100)^{2}(1)^{3} + {}^{5}C_{4}(100)(1)^{4} - {}^{5}C_{5}(1)^{5}$
= $(100)^{5} - 5(100)^{4} + 10(100)^{3} - 10(100)^{2} + 5(100) - 1$
= $1000000000 - 50000000 + 10000000 - 100000 + 500 - 1$
= $10010000500 - 500100001$
= 9509900499

Question 10:

Using Binomial Theorem, indicate which number is larger $(1.1)^{10000}$ or 1000.

Solution:

By splitting 1.1 and then applying Binomial Theorem, the first few terms of $(1.1)^{10000}$ be obtained as

$$(1.1)^{10000} = (1+0.1)^{10000}$$

= $^{10000}C_0 + ^{10000}C_1(0.1) + other positive terms$
= $1+10000 \times (0.1) + other positive terms$
= $1+1000 + other positive terms$
= $1001 + other positive terms$
> 1000

Hence, $(1.1)^{10000} > 1000$

Question 11:

Find $(a+b)^4 - (a-b)^4$. Hence, evaluate $(\sqrt{3} + \sqrt{2})^4 - (\sqrt{3} - \sqrt{2})^4$.

Solution:

Using Binomial Theorem, the expression $(a+b)^4$ and $(a-b)^4$ can be expanded as

$$(a+b)^{4} = {}^{4}C_{0}a^{4} + {}^{4}C_{1}a^{3}b + {}^{4}C_{2}a^{2}b^{2} + {}^{4}C_{3}ab^{3} + {}^{4}C_{4}b^{4}$$
$$(a-b)^{4} = {}^{4}C_{0}a^{4} - {}^{4}C_{1}a^{3}b + {}^{4}C_{2}a^{2}b^{2} - {}^{4}C_{3}ab^{3} + {}^{4}C_{4}b^{4}$$

Therefore,



$$(a+b)^{4} - (a-b)^{4} = \begin{bmatrix} ({}^{4}C_{0}a^{4} + {}^{4}C_{1}a^{3}b + {}^{4}C_{2}a^{2}b^{2} + {}^{4}C_{3}ab^{3} + {}^{4}C_{4}b^{4}) \\ - ({}^{4}C_{0}a^{4} - {}^{4}C_{1}a^{3}b + {}^{4}C_{2}a^{2}b^{2} - {}^{4}C_{3}ab^{3} + {}^{4}C_{4}b^{4}) \end{bmatrix}$$

$$= 2({}^{4}C_{1}a^{3}b + {}^{4}C_{3}ab^{3})$$

$$= 2(4a^{3}b + 4ab^{3})$$

$$= 8ab(a^{2} + b^{2})$$

Putting $a = \sqrt{3}$ and $b = \sqrt{2}$,
$$(\sqrt{3} + \sqrt{2})^{4} - (\sqrt{3} - \sqrt{2})^{4} = 8(\sqrt{3})(\sqrt{2})[(\sqrt{3})^{2} + (\sqrt{2})^{2}]$$

$$= 8\sqrt{6}[3+2]$$

$$= 40\sqrt{6}$$

Question 12:

Find $(x+1)^{6} + (x-1)^{6}$. Hence, evaluate $(\sqrt{2}+1)^{6} + (\sqrt{2}-1)^{6}$.

Solution:

Using Binomial Theorem, the expression $(x+1)^6$ and $(x-1)^6$ can be expanded as $(x+1)^6 = {}^6C_0x^6 + {}^6C_1x^5 + {}^6C_2x^4 + {}^6C_3x^3 + {}^6C_4x^2 + {}^6C_5x + {}^6C_6$ $(x-1)^6 = {}^6C_0x^6 - {}^6C_1x^5 + {}^6C_2x^4 - {}^6C_3x^3 + {}^6C_4x^2 - {}^6C_5x + {}^6C_6$

Therefore,

$$(x+1)^{6} + (x-1)^{6} = \begin{bmatrix} ({}^{6}C_{0}x^{6} + {}^{6}C_{1}x^{5} + {}^{6}C_{2}x^{4} + {}^{6}C_{3}x^{3} + {}^{6}C_{4}x^{2} + {}^{6}C_{5}x + {}^{6}C_{6}) \\ + ({}^{6}C_{0}x^{6} - {}^{6}C_{1}x^{5} + {}^{6}C_{2}x^{4} - {}^{6}C_{3}x^{3} + {}^{6}C_{4}x^{2} - {}^{6}C_{5}x + {}^{6}C_{6}) \end{bmatrix} \\ = 2 \begin{bmatrix} {}^{6}C_{0}x^{6} + {}^{6}C_{2}x^{4} + {}^{6}C_{4}x^{2} + {}^{6}C_{6} \end{bmatrix} \\ = 2 \begin{bmatrix} {}^{6}C_{0}x^{6} + {}^{6}C_{2}x^{4} + {}^{6}C_{4}x^{2} + {}^{6}C_{6} \end{bmatrix} \\ = 2 \begin{bmatrix} {}^{8}C_{1}x^{6} + {}^{1}5x^{4} + {}^{1}5x^{2} + {}^{1} \end{bmatrix}$$

Putting
$$x = \sqrt{2}$$

 $(\sqrt{2}+1)^6 + (\sqrt{2}-1)^6 = 2[(\sqrt{2})^6 + 15(\sqrt{2})^4 + 15(\sqrt{2})^2 + 1]$
 $= 2[8+15\times4+15\times2+1]$
 $= 2[8+60+30+1]$
 $= 2\times99$
 $= 198$



Question 13:

Show that $9^{n+1} - 8n - 9$ is divisible by 64, whenever *n* is a positive integer.

Solution:

To show that $9^{n+1} - 8n - 9$ is divisible by 64, we need to prove that $9^{n+1} - 8n - 9 = 64k$ where k is some natural number.

By binomial theorem,

 $(1+a)^m = {}^mC_0 + {}^mC_1a + {}^mC_2a^2 + \ldots + {}^mC_ma^m$

For a = 8 and m = n + 1, we obtain

$$(1+8)^{n+1} = {}^{n+1}C_0 + {}^{n+1}C_1(8) + {}^{n+1}C_2(8)^2 + \ldots + {}^{n+1}C_{n+1}(8)^{n+1}$$

Hence,

$$9^{n+1} = 1 + (n+1)(8) + (8)^{2} \left[{}^{n+1}C_{2} + {}^{n+1}C_{3}(8) + \dots + {}^{n+1}C_{n+1}(8)^{n+1} \right]$$

$$9^{n+1} = 9 + 8n + 64 \left[{}^{n+1}C_{2} + {}^{n+1}C_{3}(8) + \dots + {}^{n+1}C_{n+1}(8)^{n+1} \right]$$

Therefore,
$$9^{n+1} - 8n - 9 = 64k$$

Where $k = \left[{}^{n+1}C_2 + {}^{n+1}C_3(8) + \dots + {}^{n+1}C_{n+1}(8)^{n+1} \right]$ is a natural number.

Thus, $9^{n+1} - 8n - 9$ is divisible by 64, whenever *n* is a positive integer.

Question 14:

Prove that $\sum_{r=0}^{n} 3^{r} C_{r} = 4^{n}$

Solution: By Binomial theorem,

$$\sum_{r=0}^{n} {}^{n}C_{r}a^{n-r}b^{r} = (a+b)^{n}$$

By putting b = 3 and a = 1, we obtain

$$\sum_{r=0}^{n} {}^{n}C_{r} (1)^{n-r} 3^{r} = (1+3)^{n}$$
$$\sum_{r=0}^{n} 3^{r} {}^{n}C_{r} = 4^{n}$$
Hence Proved



EXERCISE 8.2

Question 1:

Find the coefficient of $x^5 in(x+3)^8$.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) in the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Assuming x^5 occurs in the expansion of $(x+3)^8$, we obtain $T_{r+1} = {}^8C_r(x)^{8-r}(3)^r$

Comparing the indices of x in x^5 in (T_{r+1}) , we obtain

$$8 - r = 5$$
$$r = 3$$

Thus, the coefficient of x^5 is ${}^{8}C_{3}(3)^{3}$

$${}^{8}C_{3}(3)^{3} = \frac{8!}{3!(5)!} \times (3)^{3}$$
$$= \frac{8 \times 7 \times 6 \times (5!)}{3 \times (2!) \times (5!)} \times 27$$
$$= 1512$$

Question 2:

Find the coefficient of a^5b^7 in $(a-2b)^{12}$

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) in the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Assuming $a^{5}b^{7}$ occurs in the expansion of $(a-2b)^{12}$, we obtain $T_{r+1} = {}^{12}C_{r}(a)^{12-r}(-2b)^{r}$ $= {}^{12}C_{r}(-2)^{r}(a)^{12-r}(b)^{r}$

Comparing the indices of a and b in a^5b^7 in (T_{r+1}) , we obtain

$$r = 7$$

Thus, the coefficient of a^5b^7 is ${}^{12}C_7(-2)^7$



$${}^{12}C_{7}(-2)^{7} = \frac{12!}{7!(5)!} \times (-2)^{7}$$
$$= \frac{12 \times 11 \times 10 \times 9 \times 8 \times (7!)}{(7!) \times (5!)} \times (-128)$$
$$= -(792) \times (128)$$
$$= -101376$$

Question 3:

Write the general term in the expansion of $(x^2 - y)^6$

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) in the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^n C_r a^{n-r} b^r$

Thus, the general term in the expansion of $(x^2 - y)^6$ is

$$T_{r+1} = {}^{6}C_{r} (x^{2})^{6-4} (-y)^{r}$$
$$= (-1)^{r} {}^{6}C_{r} (x)^{12-2r} (y)^{r}$$

Question 4:

Write the general term in expansion of $(x^2 - yx)^{12}$, $x \neq 0$

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) in the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Thus, the general term in the expansion of $(x^2 - yx)^{12}$ is

$$T_{r+1} = {}^{12}C_r (x^2)^{12-r} (-yx)^r$$

= ${}^{12}C_r (x)^{24-2r} (-1)^r (y)^r (x)^r$
= $(-1)^{r-12}C_r (x)^{24-r} (y)^r$

Question 5:

Find the 4th term in the expansion of $(x-2y)^{12}$



It is known that $(r+1)^{th}$ term, (T_{r+1}) in the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Thus, the 4th term in the expansion of $(x-2y)^{12}$ is $T_4 = T_{3+1}$ $= {}^{12}C_3(x)^{12-3}(-2y)^3$ $12! < x^3 < x^3$

$$= \frac{12!}{3!9!} (x)^{9} (-2)^{5} (y)^{9}$$
$$= \frac{12 \times 11 \times 10}{3 \times 2} \times (-8) x^{9} y^{3}$$
$$= -1760 x^{9} y^{3}$$

Question 6:

Find the 13th term in the expansion of $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}, x \neq 0$

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Thus, the 13th term in the expansion of
$$\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}, x \neq 0$$

$$T_{13} = T_{12+1}$$

$$= {}^{18}C_{12} (9x)^{18-12} \left(-\frac{1}{3\sqrt{x}}\right)^{12}$$

$$= \frac{18!}{(12!)(6!)} \times 9^6 \times (x)^6 \left(-\frac{1}{3}\right)^{12} \left(\frac{1}{\sqrt{x}}\right)^{12}$$

$$= \frac{18 \times 17 \times 16 \times 15 \times 14 \times 13 \times (12!)}{(12!) \times 6 \times 5 \times 4 \times 3 \times 2} \times (3^{12}) \times \left(\frac{1}{3^{12}}\right) \times (x^6) \times \left(\frac{1}{x^6}\right) \qquad \left[\because 9^6 = (3^2)^6 = 3^{12}\right]$$

$$= 18564$$

Question 7:

Find the middle terms in the expansion of $\left(3 - \frac{x^3}{6}\right)^7$



It is known that in the expansion of $(a+b)^n$, when *n* is odd; there are two middle terms,

namely $\left(\frac{n+1}{2}\right)^{th}$ term and $\left(\frac{n+1}{2}+1\right)^{th}$ term. $\left(2 - \frac{x^3}{2}\right)^7 = \left(7+1\right)^{th}$ where th

Therefore, the middle terms in the expansion $\left(3 - \frac{x^3}{6}\right)^7$ are $\left(\frac{7+1}{2}\right)^{th} = 4^{th}$ term and

$$\left(\frac{7+1}{2}+1\right)^{th} = 5^{th} \operatorname{term}_{T_4 = T_{3+1}}$$
$$= {}^7C_3 (3)^{7-3} \left(-\frac{x^3}{6}\right)^3$$
$$= \frac{(7!)}{(3!)(4!)} \times (3^4) \times (-1)^3 \times \left(\frac{x^9}{6^3}\right)$$
$$= -\frac{7 \times 6 \times 5 \times (4!)}{3 \times 2 \times (4!)} \times (3^4) \times \left(\frac{1}{2^3 \times 3^3}\right) \times (x^9)$$
$$= -\frac{105}{8} x^9$$

Now,

$$T_{5} = T_{4+1}$$

$$= {}^{7}C_{4} (3)^{7-4} \left(-\frac{x^{3}}{6}\right)^{4}$$

$$= \frac{7!}{(4!)(3)!} \times (3)^{3} \times \left(\frac{x^{12}}{6^{4}}\right)$$

$$= \frac{7 \cdot 6 \cdot 5 \cdot 4!}{3 \cdot 2!(4)!} \times (3^{3}) \times \left(\frac{1}{2^{4} \times 3^{4}}\right) \times (x^{12})$$

$$= \frac{35}{48} x^{12}$$

Thus, the middle terms in the expansion of $\left(3 - \frac{x^3}{6}\right)^7$ are $-\frac{105}{8}x^9$ and $\frac{35}{48}x^{12}$

Question 8:

Find the middle terms in the expansion of
$$\left(\frac{x}{3} + 9y\right)^{10}$$



It is known that in the expansion of $(a+b)^n$, when *n* is even; the middle term is $(\frac{n}{2}+1)^h$ term.

Therefore, the middle term in the expansion of $\left(\frac{x}{3} + 9y\right)^{10}$ is $\left(\frac{10}{2} + 1\right)^{th} = 6^{th}$ term

$$T_{6} = T_{5+1}$$

$$= {}^{10}C_{5} \left(\frac{x}{3}\right)^{10-5} (9y)^{5}$$

$$= \frac{(10!)}{(5!)(5!)} \times \left(\frac{x^{5}}{3^{5}}\right) \times (9^{5}) \times (y^{5})$$

$$= \frac{10 \times 9 \times 8 \times 7 \times 6 \times (5!)}{5 \times 4 \times 3 \times 2 \times (5!)} \times \left(\frac{1}{3^{5}}\right) \times (3^{10}) \times (x^{5})(y^{5})$$

$$= 252 \times 3^{5} \times x^{5} \times y^{5}$$

$$= 61236x^{5}y^{5}$$

Question 9:

In the expansion of $(1+a)^{m+n}$, prove that the coefficients of a^m and a^n are equal.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^n C_r a^{n-r} b^r$

Assuming that a^m occurs in the $\binom{r+1}{m}^m$ term of the expansion $(1+a)^{m+n}$, we obtain $T_{r+1} = {}^{m+n}C_r (1)^{m+n-r} (a)^r = {}^{m+n}C_r a^r$

Comparing the indices of a in a^m and in T_{r+1} , we obtain r = m

Therefore, the coefficient of a^m is

$$^{n+n}C_m = \frac{(m+n)!}{m!(m+n-m)!}$$

= $\frac{(m+n)!}{(m!)(n!)}$...(1)

Assuming that a^n occurs in the $(k+1)^{th}$ term of the expansion $(1+a)^{m+n}$, we obtain $T_{k+1} = {}^{m+n}C_k (1)^{m+n-k} (a)^k = {}^{m+n}C_k a^k$

Comparing the indices of *a* in a^n and in T_{k+1} , we obtain k = n



Therefore, the coefficient of a^n is

$${}^{m+n}C_n = \frac{(m+n)!}{n!(m+n-n)!}$$
$$= \frac{(m+n)!}{(m!)(n!)} \qquad \dots (2)$$

Thus from (1) and (2), it is clear that the coefficients of a^m and a^n in the expansion of $(1+a)^{m+n}$ are equal Hence proved.

Question 10:

The coefficients of the $(r-1)^{th}$, r^{th} and $(r+1)^{th}$ term in the expansion of $(x+1)^n$ are in the ratio of 1:3:5. Find *n* and *r*.

Solution:

It is known that $(k+1)^{th}$ term, (T_{k+1}) term of the expansion $(a+b)^n$ is given by $T_{k+1} = {}^{n}C_k a^{n-k}b^k$

Hence, $(r-1)^{th}$ term in the expansion of $(x+1)^n$ is $T_{r-1} = {}^n C_{r-2} (x)^{n-(r-2)} (1)^{(r-2)}$ $= {}^n C_{r-2} x^{n-r+2}$

 $(r+1)^{th}$ term in the expansion of $(x+1)^{th}$ is

$$T_{r+1} = {^{n}C_{r}(x)^{n-r}(1)^{r}} = {^{n}C_{r}x^{n-r}}$$

 r^{th} term in the expansion of $(x+1)^n$ is

$$T_{r} = {}^{n}C_{r-1}(x)^{n-(r-1)}(1)^{(r-1)}$$
$$= {}^{n}C_{r-1}x^{n-r+1}$$

Therefore, coefficients of the $(r-1)^{th}$, r^{th} and $(r+1)^{th}$ term in the expansion of $(x+1)^{n}$ are ${}^{n}C_{r-2}$, ${}^{n}C_{r-1}$ and ${}^{n}C_{r}$ respectively.

Since these coefficients are in the ratio of 1:3:5, we obtain



$$\frac{{}^{n}C_{r-2}}{{}^{n}C_{r-1}} = \frac{1}{3} \text{ and } \frac{{}^{n}C_{r-1}}{{}^{n}C_{r}} = \frac{3}{5}$$

$$\frac{{}^{n}C_{r-2}}{{}^{n}C_{r-1}} = \frac{n!}{(r-2)!(n-r+2)!} \times \frac{(r-1)!(n-r+1)!}{n!}$$

$$\frac{1}{3} = \frac{(r-1).(r-2)!(n-r+1)!}{(r-2)!(n-r+2)!(n-r+1)!}$$

$$\frac{1}{3} = \frac{r-1}{n-r+2}$$

$$n-r+2 = 3r-3$$

$$n-4r+5 = 0$$

$$n = 4r-5 \qquad \dots(1)$$

$$\frac{{}^{n}C_{r-1}}{{}^{n}C_{r}} = \frac{n!}{(r-1)!(n-r+1)!} \times \frac{r!(n-r)!}{n!}$$

$$\frac{3}{5} = \frac{r.(r-1)!(n-r+1)!(n-r)!}{(r-1)!(n-r+1)!(n-r)!}$$

$$\frac{3}{5} = \frac{r}{n-r+1}$$

$$3n-3r+3 = 5r$$

$$3n-8r+3 = 0 \qquad \dots(2)$$

From (1) and (2), we obtain

$$3(4r-5)-8r+3 = 0$$

$$12r-15-8r+3 = 0$$

$$4r-12 = 0$$

$$r = 3$$

Putting the value of r in (1), we obtain n

$$n = 4 \times 3 - 5$$
$$= 12 - 5$$
$$= 7$$

Thus, n = 7 and r = 3



Question 11:

Prove that the coefficient of x^n in the expansion of $(1+x)^{2n}$ is twice the coefficient of x^n in the expansion of $(1-x)^{2n-1}$.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^n C_r a^{n-r} b^r$

Assuming that x^n occurs in the $(r+1)^{th}$ term of the expansion $(1+x)^{2n}$, we obtain $T_{r+1} = {}^{2n}C_r(1)^{2n-r}(x)^r$ $= {}^{2n}C_rx^r$

Comparing the indices of x in x^n and in T_{r+1} , we obtain r = n

Therefore, the coefficient of x^n in the expansion of $(1+x)^{2n}$

$${}^{2n}C_{n} = \frac{(2n)!}{n!(2n-n)!}$$
$$= \frac{(2n)!}{(n!)(n!)}$$
$$= \frac{(2n)!}{(n!)^{2}} \qquad \dots (1)$$

Assuming that x^n occurs in the $(k+1)^{th}$ term of the expansion $(1+x)^{2n-1}$, we obtain $T_{k+1} = {}^{2n}C_k (1)^{2n-1-k} (x)^k$ $= {}^{2n}C_k x^k$

Comparing the indices of x in x^n and in T_{k+1} , we obtain k = n

Therefore, the coefficient of x^n in the expansion of $(1+x)^{2n-1}$



$${}^{2n-1}C_n = \frac{(2n-1)!}{n!(2n-1-n)!}$$

$$= \frac{(2n-1)!}{n!(n-1)!}$$

$$= \frac{2n \cdot (2n-1)!}{2n \cdot n!(n-1)!}$$

$$= \frac{(2n)!}{2 \cdot (n!)(n!)}$$

$$= \frac{1}{2} \left[\frac{(2n)!}{(n!)^2} \right] \dots (2)$$

From (1) and (2), we can observe that

$${}^{2n-1}C_n = \frac{1}{2} \left({}^{2n}C_n \right)$$
$${}^{2n}C_n = 2 \left({}^{2n-1}C_n \right)$$

Therefore, the coefficient of x^n in the expansion of $(1+x)^{2n}$ is twice the coefficient of x^n in the expansion of $(1-x)^{2n-1}$. Hence Proved

Question 12:

Find a positive value of *m* for which the coefficient of x^2 in the expansion $(1+x)^m$ is 6.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^n$ is given by $T_{r+1} = {}^n C_r a^{n-r} b^r$

Assuming that x^2 occurs in the $(r+1)^{th}$ term of the expansion $(1+x)^m$, we obtain $T_{r+1} = {}^{m}C_r(1)^{m-r}(x)^r = {}^{m}C_r(x)^r$

Comparing the indices of x in x^2 in (T_{r+1}) , we obtain r = 2

Therefore, the coefficient of x^2 is ${}^{m}C_2$.



$${}^{m}C_{2} = 6$$

$$\frac{m!}{2!(m-2)!} = 6$$

$$\frac{m(m-1) \times (m-2)!}{2 \times (m-2)!} = 6$$

$$m(m-1) = 12$$

$$m^{2} - m - 12 = 0$$

$$m^{2} - 4m + 3m - 12 = 0$$

$$m(m-4) + 3(m-4) = 0$$

$$(m-4)(m+3) = 0$$

$$m = 4 \text{ or } m = -3$$

Thus, 4 is the positive value of *m* for which the coefficient of x^2 in the expansion $(1+x)^m$ is 6.



MISCELLANEOUS EXERCISE

Question 1:

Find a, b and n in the expansion of $(a+b)^n$ if the first three terms of the expansion are 729, 7290 and 30375 respectively.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^{n}$ is given by $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

The first three terms of expansion are 729, 7290 and 30375 respectively.

$$T_{1} = {}^{n}C_{0}a^{n-0}b^{0}$$

$$= a^{n}$$

$$= 729 \qquad \dots(1)$$

$$T_{2} = {}^{n}C_{1}a^{n-1}b^{1}$$

$$= na^{n-1}b$$

$$= 7290 \qquad \dots(2)$$

$$T_{3} = {}^{n}C_{1}a^{n-2}b^{2}$$

$$= \frac{n(n-1)}{2}a^{n-2}b^{2}$$

$$= 30375 \qquad (3)$$

Dividing (2) by (1), we obtain

$$\frac{na^{n-1}b}{a^n} = \frac{7290}{729}$$
$$\frac{nb}{a} = 10 \qquad \dots (4)$$

Dividing (3) by (2), we obtain



$$\frac{n(n-1)a^{n-2}b^2}{2na^{n-1}b} = \frac{30375}{7290}$$
$$\frac{(n-1)b}{2a} = \frac{30375}{7290}$$
$$\frac{(n-1)b}{a} = \frac{30375 \times 2}{7290}$$
$$\frac{nb}{a} - \frac{b}{a} = \frac{25}{3}$$
$$10 - \frac{b}{a} = \frac{25}{3} \qquad \dots \text{(from 4)}$$

$$\frac{b}{a} = 10 - \frac{25}{3}$$
$$\frac{b}{a} = \frac{5}{3} \qquad \dots (5)$$

Putting
$$\frac{b}{a} = \frac{5}{3}$$
 in (4), we obtain

$$n \times \frac{5}{3} = 10$$
$$n = 6$$

Substituting n = 6 in (1), we obtain

$$a^{6} = 729$$
$$a = \sqrt[6]{729}$$
$$= 3$$

From (5), we obtain

$$\frac{b}{3} = \frac{5}{3}$$
$$b = 5$$

Therefore, a = 3, b = 5, n = 6

Question 2:

Find *a* if the coefficients of x^2 and x^3 in the expansion of $(3 + ax)^9$ are equal.

Solution:

It is known that $(r+1)^{th}$ term, (T_{r+1}) the binomial expression of $(a+b)^{n}$ is given by



 $T_{r+1} = {}^{n}C_{r}a^{n-r}b^{r}$

Assuming that x^2 occurs in the $(r+1)^{th}$ term of the expansion $(3+ax)^9$, we obtain $T_{r+1} = {}^9C_r (3)^{9-r} (ax)^r$ $= {}^9C_r (3)^{9-r} a^r x^r$

Comparing the indices of x in x^2 and in (T_{r+1}) , we obtain r = 2

Thus, the coefficient of x^2 is

$${}^{9}C_{2}(3)^{9-2} a^{2} = \frac{9!}{2!7!}(3)^{7} a^{2}$$
$$= 36(3)^{7} a^{2}$$

Assuming that x^3 occurs in the $(k+1)^{th}$ term of the expansion $(3+ax)^9$, we obtain

$$T_{k+1} = {}^{9}C_{k} (3)^{9-k} (ax)^{k}$$
$$= {}^{9}C_{k} (3)^{9-k} a^{k} x^{k}$$

Comparing the indices in x^2 and x^3 in (T_{r+1}) , we obtain k=3

Thus, the coefficient of x^3 is

$${}^{9}C_{3}(3)^{9-3} a^{3} = \frac{9!}{3!6!}(3)^{6} a^{3}$$
$$= 84(3)^{6} a^{3}$$

It is given that the coefficient of x^2 and x^3 are equal.

$$84(3)^{6} a^{3} = 36(3)^{7} a^{2}$$

$$84a = 36 \times 3$$

$$a = \frac{36 \times 3}{84}$$

$$= \frac{9}{7}$$
guired value of $a = \frac{9}{7}$

Hence, the required value of u^{-1}

Question 3:

Find the coefficient of x^5 in the product $(1+2x)^6(1-x)^7$ using binomial theorem.



Using binomial theorem, $(1+2x)^6$ and $(1-x)^7$ can be expanded as $(1+2x)^6 = {}^6C_0 + {}^6C_1(2x) + {}^6C_2(2x)^2 + {}^6C_3(2x)^3 + {}^6C_4(2x)^4 + {}^6C_5(2x)^5 + {}^6C_6(2x)^6$ $= 1+6(2x)+15(2x)^2+20(2x)^3+15(2x)^4+6(2x)^5+(2x)^6$ $= 1+12x+60x^2+160x^3+240x^4+192x^5+64x^6$

$$(1-x)^{7} = {}^{7}C_{0} - {}^{7}C_{1}(x) + {}^{7}C_{2}(x)^{2} - {}^{7}C_{3}(x)^{3} + {}^{7}C_{4}(x)^{4} - {}^{7}C_{5}(x)^{5} + {}^{7}C_{6}(x)^{6} - {}^{7}C_{7}(x)^{7}$$

= 1 - 7x + 21x² - 35x³ + 35x⁴ - 21x⁵ + 7x⁶ - x⁷

Therefore,

$$(1+2x)^{6}(1-x)^{7} = (1+12x+60x^{2}+160x^{3}+240x^{4}+192x^{5}+64x^{6})(1-7x+21x^{2}-35x^{3}+35x^{4}-21x^{5}+7x^{6}-x^{7})$$

= 1(-21x^{5})+(12x)(35x^{4})+(60x^{2})(-35x^{3})+(160x^{3})(21x^{2})+(240x^{4})(-7x)+(192x^{5})(1)
= 171x⁵

Thus, the coefficient of x^5 in the product $(1+2x)^6(1-x)^7$ is 171.

Question 4:

If a and b are distinct integers, prove that a-b is a factor of $a^n - b^n$ whenever n is a positive integer.

[Hint: write $a^n = (a-b+b)^n$ and expand]

Solution:

In order to prove that a-b is a factor of $a^n - b^n$, we have to prove that $a^n - b^n = k(a-b)$ where k is some natural number.

We can write *a* as

$$a = a - b + b$$

$$a^{n} = (a - b + b)^{n}$$

$$a^{n} = [(a - b) + b]^{n}$$

$$a^{n} = {nC_{0}(a - b)^{n} + {nC_{1}(a - b)^{n-1}b + \dots + {nC_{n-1}(a - b)b^{n-1} + {nC_{n}b^{n}}}}$$

$$a^{n} = (a - b)^{n} + {nC_{1}(a - b)^{n-1}b + \dots + {nC_{n-1}(a - b)b^{n-1} + b^{n}}}$$

$$a^{n} - b^{n} = (a - b)[(a - b)^{n-1} + {nC_{1}(a - b)^{n-2}b + \dots + {nC_{n-1}b^{n-1}}}]$$

$$a^{n} - b^{n} = k(a - b)$$
where $k = [(a - b)^{n-1} + {nC_{1}(a - b)^{n-2}b + \dots + {nC_{n-1}b^{n-1}}]$



This shows that a-b is a factor of $a^n - b^n$ whenever *n* is a positive integer. Hence proved.

Question 5:

Evaluate
$$\left(\sqrt{3}+\sqrt{2}\right)^6 - \left(\sqrt{3}-\sqrt{2}\right)^6$$

Solution:

Using binomial theorem,

$$(a+b)^{6} = {}^{6}C_{0}a^{6} + {}^{6}C_{1}a^{5}b + {}^{6}C_{2}a^{4}b^{2} + {}^{6}C_{3}a^{3}b^{3} + {}^{6}C_{4}a^{2}b^{4} + {}^{6}C_{5}ab^{5} + {}^{6}C_{6}b^{6}$$
$$= a^{6} + 6a^{5}b + 15a^{4}b^{2} + 20a^{3}b^{3} + 15a^{2}b^{4} + 6ab^{5} + b^{6}$$

$$(a-b)^{6} = {}^{6}C_{0}a^{6} - {}^{6}C_{1}a^{5}b + {}^{6}C_{2}a^{4}b^{2} - {}^{6}C_{3}a^{3}b^{3} + {}^{6}C_{4}a^{2}b^{4} - {}^{6}C_{5}ab^{5} + {}^{6}C_{6}b^{6}$$
$$= a^{6} - 6a^{5}b + 15a^{4}b^{2} - 20a^{3}b^{3} + 15a^{2}b^{4} - 6ab^{5} + b^{6}$$

Therefore,

$$(a+b)^{6} - (a-b)^{6} = 2 \left[6a^{5}b + 20a^{3}b^{3} + 6ab^{5} \right]$$

Putting $a = \sqrt{3}$ and $b = \sqrt{2}$, we obtain,

$$\left(\sqrt{3} + \sqrt{2}\right)^6 - \left(\sqrt{3} - \sqrt{2}\right)^6 = 2 \left[6\left(\sqrt{3}\right)^5 \left(\sqrt{2}\right) + 20\left(\sqrt{3}\right)^3 \left(\sqrt{2}\right)^3 + 6\left(\sqrt{3}\right) \left(\sqrt{2}\right)^5\right]$$

= 2 \left[54\sqrt{6} + 120\sqrt{6} + 24\sqrt{6} \right]
= 2 \times 198\sqrt{6}
= 396\sqrt{6}

Question 6:

Find the value of $(a^2 + \sqrt{a^2 - 1})^4 + (a^2 - \sqrt{a^2 - 1})^4$

Solution:

Using binomial theorem,

$$(x+y)^{4} = {}^{4}C_{0}x^{4} + {}^{4}C_{1}x^{3}y + {}^{4}C_{2}x^{2}y^{2} + {}^{4}C_{3}xy^{3} + {}^{4}C_{4}y^{4}$$
$$= x^{4} + 4x^{3}y + 6x^{2}y^{2} + 4xy^{3} + y^{4}$$

$$(x-y)^{4} = {}^{4}C_{0}x^{4} - {}^{4}C_{1}x^{3}y + {}^{4}C_{2}x^{2}y^{2} - {}^{4}C_{3}xy^{3} + {}^{4}C_{4}y^{4}$$
$$= x^{4} - 4x^{3}y + 6x^{2}y^{2} - 4xy^{3} + y^{4}$$



Therefore,

$$(x+y)^4 + (x-y)^4 = 2(x^4 + 6x^2y^2 + y^4)$$

Putting $x = a^2$ and $y = \sqrt{a^2 - 1}$, we obtain

$$\left(a^{2} + \sqrt{a^{2} - 1}\right)^{4} + \left(a^{2} - \sqrt{a^{2} - 1}\right)^{4} = 2\left[\left(a^{2}\right)^{4} + 6\left(a^{2}\right)^{2}\left(\sqrt{a^{2} - 1}\right)^{2} + \left(\sqrt{a^{2} - 1}\right)^{4}\right]$$

$$= 2\left[a^{8} + 6a^{4}\left(a^{2} - 1\right) + \left(a^{2} - 1\right)^{2}\right]$$

$$= 2\left[a^{8} + 6a^{6} - 6a^{4} + a^{4} - 2a^{2} + 1\right]$$

$$= 2\left[a^{8} + 6a^{6} - 5a^{4} - 2a^{2} + 1\right]$$

$$= 2a^{8} + 12a^{6} - 10a^{4} - 4a^{2} + 2$$

Question 7:

Find an approximation of $(0.99)^5$ using the first three terms of its expansion.

Solution:

We can express 0.99 as the difference of two numbers whose powers are easier to calculate.

Hence, 0.99 = 1 - 0.01

Therefore,

$$(0.99)^{5} = (1 - 0.01)^{5}$$

= ${}^{5}C_{0}(1)^{5} - {}^{5}C_{1}(1)^{4}(0.01) + {}^{5}C_{2}(1)^{3}(0.01)^{2}$
= $1 - 5(0.01) + 10(0.01)^{2}$
= $1 - 0.05 + 0.001$
= $1.001 - 0.05$
= 0.951

Hence, the approximation of $(0.99)^5$ is 0.951

Question 8:

Find *n*, if the ratio of the fifth term from the beginning to the fifth term from the end in the

expansion of
$$\left(\sqrt[4]{2} + \frac{1}{\sqrt[4]{3}}\right)^n$$
 is $\sqrt{6}:1$



In the expansion,

$$(a+b)^{n} = {}^{n}C_{0}a^{n} + {}^{n}C_{1}a^{n-1}b + {}^{n}C_{2}a^{n-2}b^{2} + \dots + {}^{n}C_{n-1}ab^{n-1} + {}^{n}C_{n}b^{n}$$

Fifth term from the beginning is ${}^{n}C_{4}a^{n-4}b^{4}$ Fifth term from the end is ${}^{n}C_{4}a^{4}b^{n-4}$

Therefore, in the expansion of $\left(\sqrt[4]{2} + \frac{1}{\sqrt[4]{3}}\right)^n$;

fifth term from the beginning is
$${}^{n}C_{4}\left(\frac{4}{\sqrt{2}}\right)^{n-4}\left(\frac{1}{\sqrt[4]{3}}\right)^{4}$$
 and

fifth term from the end is ${}^{n}C_{n-4}\left(\sqrt[4]{2}\right)^{4}\left(\frac{1}{\sqrt[4]{3}}\right)^{n-4}$

$${}^{n}C_{4}\left(\sqrt[4]{2}\right)^{n-4}\left(\frac{1}{\sqrt[4]{3}}\right)^{4} = {}^{n}C_{4}\frac{\left(\sqrt[4]{2}\right)^{n}}{\left(\sqrt[4]{2}\right)^{4}} \cdot \frac{1}{3}$$
$$= \frac{n!}{6 \cdot 4!(n-4)!} \left(\sqrt[4]{2}\right)^{n} \qquad \dots (1)$$

$${}^{n}C_{n-4} \left(\sqrt[4]{2}\right)^{4} \left(\frac{1}{\sqrt[4]{3}}\right)^{n-4} = {}^{n}C_{n-4} \frac{\left(\sqrt[4]{3}\right)^{4}}{\left(\sqrt[4]{3}\right)^{n}}$$
$$= {}^{n}C_{n-4} \cdot 2 \cdot \frac{3}{\left(\sqrt[4]{3}\right)^{n}}$$
$$= \frac{6n!}{(n-4)!4!} \cdot \frac{1}{\left(\sqrt[4]{3}\right)^{n}} \qquad \dots (2)$$

It is given that the ratio of of the fifth term from the beginning to the fifth term from the end

in the expansion of $\left(\sqrt[4]{2} + \frac{1}{\sqrt[4]{3}}\right)^n$ is $\sqrt{6}:1$

Therefore,



$$\left[\frac{n!}{6\cdot 4!(n-4)!} \left(\frac{4}{2}\right)^{n}\right] : \left[\frac{6n!}{(n-4)!4!} \cdot \frac{1}{\left(\frac{4}{\sqrt{3}}\right)^{n}}\right] = \sqrt{6} : 1$$
$$\frac{\left(\frac{4}{\sqrt{2}}\right)^{n}}{6} : \frac{6}{\left(\frac{4}{\sqrt{3}}\right)^{n}} = \sqrt{6} : 1$$
$$\frac{\left(\frac{4}{\sqrt{2}}\right)^{n}}{6} \times \frac{\left(\frac{4}{\sqrt{3}}\right)^{n}}{6} = \sqrt{6}$$
$$\left(\frac{4}{\sqrt{6}}\right)^{n} = 36\sqrt{6}$$
$$\left(\frac{6}{\sqrt{6}}\right)^{n} = 36\sqrt{6}$$
$$\left(\frac{6}{\sqrt{6}}\right)^{n} = \frac{5}{2}$$
$$n = 4 \times \frac{5}{2}$$
$$= 10$$

Thus the value of n = 10.

Question 9:

Expand using binomial theorem $\left(1+\frac{x}{2}-\frac{2}{x}\right)^4, x \neq 0$

Solution:

$$\left(1 + \frac{x}{2} - \frac{2}{x}\right)^4 = {}^nC_0 \left(1 + \frac{x}{2}\right)^4 - {}^nC_1 \left(1 + \frac{x}{2}\right)^3 \left(\frac{2}{x}\right) + {}^nC_2 \left(1 + \frac{x}{2}\right)^2 \left(\frac{2}{x}\right)^2 - {}^nC_3 \left(1 + \frac{x}{2}\right) \left(\frac{2}{x}\right)^3 + {}^nC_4 \left(\frac{2}{x}\right)^4$$

$$= \left(1 + \frac{x}{2}\right)^4 - 4\left(1 + \frac{x}{2}\right)^3 \left(\frac{2}{x}\right) + 6\left(1 + x + \frac{x^2}{4}\right) \left(\frac{4}{x^2}\right) - 4\left(1 + \frac{x}{2}\right) \left(\frac{8}{x^3}\right) + \frac{16}{x^4}$$

$$= \left(1 + \frac{x}{2}\right)^4 - \frac{8}{x} \left(1 + \frac{x}{2}\right)^3 + \frac{24}{x^2} + \frac{24}{x} + 6 - \frac{32}{x^3} - \frac{16}{x^2} + \frac{16}{x^4}$$

$$= \left(1 + \frac{x}{2}\right)^4 - \frac{8}{x} \left(1 + \frac{x}{2}\right)^3 + \frac{8}{x^2} + \frac{24}{x} + 6 - \frac{32}{x^3} + \frac{16}{x^4}$$

$$\dots (1)$$



Again, by using binomial theorem, we obtain

$$\left(1+\frac{x}{2}\right)^{4} = {}^{4}C_{0}\left(1\right)^{4} + {}^{4}C_{1}\left(1\right)^{3}\left(\frac{x}{2}\right) + {}^{4}C_{2}\left(1\right)^{2}\left(\frac{x}{2}\right)^{2} + {}^{4}C_{3}\left(1\right)\left(\frac{x}{2}\right)^{3} + {}^{4}C_{4}\left(\frac{x}{2}\right)^{4}$$
$$= 1+4\times\frac{x}{2}+6\times\frac{x^{4}}{4}+4\times\frac{x^{3}}{8}+\frac{x^{4}}{16}$$
$$= 1+2x+\frac{3x^{2}}{2}+\frac{x^{3}}{2}+\frac{x^{4}}{16} \qquad \dots(2)$$

And

$$\left(1+\frac{x}{2}\right)^{3} = {}^{3}C_{0}\left(1\right)^{3} + {}^{3}C_{1}\left(1\right)^{2}\left(\frac{x}{2}\right) + {}^{3}C_{2}\left(1\right)\left(\frac{x}{2}\right)^{2} + {}^{3}C_{3}\left(\frac{x}{2}\right)^{3}$$
$$= 1+\frac{3x}{2}+\frac{3x^{2}}{4}+\frac{x^{3}}{8} \qquad \dots(3)$$

From (1), (2) and (3), we obtain

$$\left[\left(1+\frac{x}{2}\right)-\frac{2}{x}\right]^{4} = 1+2x+\frac{3x^{2}}{2}+\frac{x^{3}}{2}+\frac{x^{4}}{16}-\frac{8}{x}\left(1+\frac{3x}{2}+\frac{3x^{2}}{4}+\frac{x^{3}}{8}\right)+\frac{8}{x^{2}}+\frac{24}{x}+6-\frac{32}{x^{3}}+\frac{16}{x^{4}}\right]$$
$$= 1+2x+\frac{3x^{2}}{2}+\frac{x^{3}}{2}+\frac{x^{4}}{16}-\frac{8}{x}-12-6x-x^{2}+\frac{8}{x^{2}}+\frac{24}{x}+6-\frac{32}{x^{3}}+\frac{16}{x^{4}}$$
$$= \frac{16}{x}+\frac{8}{x^{2}}-\frac{32}{x^{3}}+\frac{16}{x^{4}}-4x+\frac{x^{2}}{2}+\frac{x^{3}}{2}+\frac{x^{4}}{16}-5$$

Question 10:

Find the expansion of $(3x^2 - 2ax + 3a^2)^3$

Solution:

Using binomial theorem, the expansion of
$$(3x^2 - 2ax + 3a^2)^3$$
 is given by

$$\begin{bmatrix} (3x^2 - 2ax) + 3a^2 \end{bmatrix}^3 = {}^{3}C_0 (3x^2 - 2ax)^3 + {}^{3}C_1 (3x^2 - 2ax)^2 (3a^2) + {}^{3}C_2 (3x^2 - 2ax) (3a^2)^2 + {}^{3}C_3 (3a^2)^3$$

$$= (3x^2 - 2ax)^3 + 3(9x^4 - 12ax^3 + 4a^2x^2) (3a^2) + 3(3x^2 - 2ax) (9a^4) + 27a^6$$

$$= (3x^2 - 2ax)^3 + 81a^2x^4 - 108a^3x^3 + 36a^4x^2 + 81a^4x^2 - 54a^5x + 27a^6$$

$$= (3x^2 - 2ax)^3 + 81a^2x^4 - 108a^3x^3 + 117a^4x^2 - 54a^5x + 27a^6$$
...(1)

Again, by using binomial theorem, we obtain



$$(3x^{2} - 2ax)^{3} = {}^{3}C_{0}(3x^{2})^{3} - {}^{3}C_{1}(3x^{2})^{2}(2ax) + {}^{3}C_{2}(3x^{2})(2ax)^{2} + {}^{3}C_{3}(2ax)^{3}$$

= 27x⁶ - 3(9x⁴)(2ax) + 3(3x²)(4a²x²) - 8a³x³
= 27x⁶ - 54ax⁵ + 36a²x⁴ - 8a³x³ ...(2)

From (1) and (2), we obtain

$$(3x^{2} - 2ax + 3a^{2})^{3} = 27x^{6} - 54ax^{5} + 36a^{2}x^{4} - 8a^{3}x^{3} + 81a^{2}x^{4} - 108a^{3}x^{3} + 117a^{4}x^{2} - 54a^{5}x + 27a^{6}$$
$$= 27x^{6} - 54ax^{5} + 117a^{2}x^{4} - 116a^{3}x^{3} + 117a^{4}x^{2} - 54a^{5}x + 27a^{6}$$





When you learn math in an interesting way, you never forget.



25 Million

Math classes & counting

100K+

Students learning Math the right way

20+ Countries

Present across USA, UK, Singapore, India, UAE & more.

Why choose Cuemath?

"Cuemath is a valuable addition to our family. We love solving puzzle cards. My daughter is now visualizing maths and solving problems effectively!" "Cuemath is great because my son has a one-on-one interaction with the teacher. The instructor has developed his confidence and I can see progress in his work. One-on-one interaction is perfect and a great bonus." "I appreciate the effort that miss Nitya puts in to help my daughter understand the best methods and to explain why she got a problem incorrect. She is extremely patient and generous with Miranda."

- Gary Schwartz

- Kirk Riley

- Barbara Cabrera

Get the Cuemath advantage

Book a FREE trial class